


# OPTIMAL INSTRUMENT

## Difficulty–Baseline

Instructions: Please circle the level of difficulty you have for each activity today.	Able to do without any difficulty	Able to do with little difficulty	Able to do with moderate difficulty	Able to do with much difficulty	Unable to do	Not applicable
1. Lying flat	1	2	3	4	5	9
2. Rolling over	1	2	3	4	5	9
3. Moving–lying to sitting	1	2	3	4	5	9
4. Sitting	1	2	3	4	5	9
5. Squatting	1	2	3	4	5	9
6. Bending/stooping	1	2	3	4	5	9
7. Balancing	1	2	3	4	5	9
8. Kneeling	1	2	3	4	5	9
9. Walking–short distance	1	2	3	4	5	9
10. Walking–long distance	1	2	3	4	5	9
11. Walking–outdoors	1	2	3	4	5	9
12. Climbing stairs	1	2	3	4	5	9
13. Hopping	1	2	3	4	5	9
14. Jumping	1	2	3	4	5	9
15. Running	1	2	3	4	5	9
16. Pushing	1	2	3	4	5	9
17. Pulling	1	2	3	4	5	9
18. Reaching	1	2	3	4	5	9
19. Grasping	1	2	3	4	5	9
20. Lifting	1	2	3	4	5	9
21. Carrying	1	2	3	4	5	9

22. Thinking about all of the activities you would like to do, please mark an “X” at the point on the line that best describes your overall level of difficulty with these activities today.



23. From the above list, choose the 3 activities you would most like to be able to do without any difficulty (for example, if you would most like to be able to *climb stairs*, *kneel*, and *hop* without any difficulty, you would choose: 1. 12 2. 8 3. 13 )

1. \_\_\_\_ 2. \_\_\_\_ 3. \_\_\_\_


© 2005, 2006 American Physical Therapy Association. All rights reserved. No part of this instrument may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, or otherwise without prior permission of the American Physical Therapy Association. Contact [permissions@apta.org](mailto:permissions@apta.org) or visit [www.apta.org/publications](http://www.apta.org/publications).

Adapted/revised in July 2005 and August 2006 with permission of APTA from Guccione AA, Mielenz TJ, De Vellis RF, et al. Development and testing of a self-report instrument to measure actions: Outpatient Physical Therapy Improvement in Movement Assessment Log (OPTIMAL). *Phys Ther.* 2005;85:515-530.

### Confidence–Baseline

Instructions: Please circle the level of confidence you have for doing each activity today.	Fully confident in my ability to perform	Very confident	Moderate confidence	Some confidence	Not confident in my ability to perform	Not applicable
1. Lying flat	1	2	3	4	5	9
2. Rolling over	1	2	3	4	5	9
3. Moving–lying to sitting	1	2	3	4	5	9
4. Sitting	1	2	3	4	5	9
5. Squatting	1	2	3	4	5	9
6. Bending/stooping	1	2	3	4	5	9
7. Balancing	1	2	3	4	5	9
8. Kneeling	1	2	3	4	5	9
9. Walking–short distance	1	2	3	4	5	9
10. Walking–long distance	1	2	3	4	5	9
11. Walking–outdoors	1	2	3	4	5	9
12. Climbing stairs	1	2	3	4	5	9
13. Hopping	1	2	3	4	5	9
14. Jumping	1	2	3	4	5	9
15. Running	1	2	3	4	5	9
16. Pushing	1	2	3	4	5	9
17. Pulling	1	2	3	4	5	9
18. Reaching	1	2	3	4	5	9
19. Grasping	1	2	3	4	5	9
20. Lifting	1	2	3	4	5	9
21. Carrying	1	2	3	4	5	9

22. Thinking about all the activities you like to do, please mark an “X” at the point on the line that best describes your overall level of confidence in performing these activities today:



© 2005, 2006 American Physical Therapy Association. All rights reserved. No part of this instrument may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, or otherwise without prior permission of the American Physical Therapy Association. Contact [permissions@apta.org](mailto:permissions@apta.org) or visit [www.apta.org/publications](http://www.apta.org/publications).

Adapted/revised in July 2005 and August 2006 with permission of APTA from Guccione AA, Mielenz TJ, De Vellis RF, et al. Development and testing of a self-report instrument to measure actions: Outpatient Physical Therapy Improvement in Movement Assessment Log (OPTIMAL). *Phys Ther.* 2005;85:515-530.